

STUDY Guide FINAL
NO CALC

1. If $y = (x^3 + 1)^2$, then $\frac{dy}{dx} =$

(A) $(3x^2)^2$

(B) $2(x^3 + 1)$

(C) $2(3x^2 + 1)$

(D) $3x^2(x^3 + 1)$

(E) $6x^2(x^3 + 1)$

2. $\int_0^1 e^{-4x} dx =$

(A) $\frac{-e^{-4}}{4}$

(B) $-4e^{-4}$

(C) $e^{-4} - 1$

(D) $\frac{1}{4} - \frac{e^{-4}}{4}$

(E) $4 - 4e^{-4}$

~~Study Guide FINAL~~~~NO CALC~~

3. For $x \geq 0$, the horizontal line $y = 2$ is an asymptote for the graph of the function f . Which of the following statements must be true?

- (A) $f(0) = 2$
- (B) $f(x) \neq 2$ for all $x \geq 0$
- (C) $f(2)$ is undefined.
- (D) $\lim_{x \rightarrow 2} f(x) = \infty$
- (E) $\lim_{x \rightarrow \infty} f(x) = 2$

4. If $y = \frac{2x + 3}{3x + 2}$, then $\frac{dy}{dx} =$


- (A) $\frac{12x + 13}{(3x + 2)^2}$
 - (B) $\frac{12x - 13}{(3x + 2)^2}$
 - (C) $\frac{5}{(3x + 2)^2}$
 - (D) $\frac{-5}{(3x + 2)^2}$
 - (E) $\frac{2}{3}$
-

5. $\int_0^{\pi/4} \sin x \, dx =$

- (A) $-\frac{\sqrt{2}}{2}$ (B) $\frac{\sqrt{2}}{2}$ (C) $-\frac{\sqrt{2}}{2} - 1$ (D) $-\frac{\sqrt{2}}{2} + 1$ (E) $\frac{\sqrt{2}}{2} - 1$
-

6. $\lim_{x \rightarrow \infty} \frac{x^3 - 2x^2 + 3x - 4}{4x^3 - 3x^2 + 2x - 1} =$

- (A) 4 (B) 1 (C) $\frac{1}{4}$ (D) 0 (E) -1
-

Graph of f'

7. The graph of f' , the derivative of the function f , is shown above. Which of the following statements is true about f ?
- (A) f is decreasing for $-1 \leq x \leq 1$.
 - (B) f is increasing for $-2 \leq x \leq 0$.
 - (C) f is increasing for $1 \leq x \leq 2$.
 - (D) f has a local minimum at $x = 0$.
 - (E) f is not differentiable at $x = -1$ and $x = 1$.
-

8. $\int x^2 \cos(x^3) dx =$

(A) $-\frac{1}{3} \sin(x^3) + C$

(B) $\frac{1}{3} \sin(x^3) + C$

(C) $-\frac{x^3}{3} \sin(x^3) + C$

(D) $\frac{x^3}{3} \sin(x^3) + C$

(E) $\frac{x^3}{3} \sin\left(\frac{x^4}{4}\right) + C$

9. If $f(x) = \ln(x + 4 + e^{-3x})$, then $f'(0)$ is

(A) $-\frac{2}{5}$

(B) $\frac{1}{5}$


(C) $\frac{1}{4}$

(D) $\frac{2}{5}$


(E) nonexistent

10. The function f has the property that $f(x)$, $f'(x)$, and $f''(x)$ are negative for all real values x . Which of the following could be the graph of f ?


(A)




(B)




(C)



(D)



(E)




(11) Using the substitution $u = 2x + 1$, $\int_0^2 \sqrt{2x + 1} \, dx$ is equivalent to

- (A) $\frac{1}{2} \int_{-1/2}^{1/2} \sqrt{u} \, du$ (B) $\frac{1}{2} \int_0^2 \sqrt{u} \, du$ (C) $\frac{1}{2} \int_1^5 \sqrt{u} \, du$ (D) $\int_0^2 \sqrt{u} \, du$ (E) $\int_1^5 \sqrt{u} \, du$

12. The rate of change of the volume, V , of water in a tank with respect to time, t , is directly proportional to the square root of the volume. Which of the following is a differential equation that describes this relationship?

- (A) $V(t) = k\sqrt{t}$
(B) $V(t) = k\sqrt{V}$
(C) $\frac{dV}{dt} = k\sqrt{t}$
(D) $\frac{dV}{dt} = \frac{k}{\sqrt{V}}$
(E) $\frac{dV}{dt} = k\sqrt{V}$
-

Graph of f

13. The graph of a function f is shown above. At which value of x is f continuous, but not differentiable?

- (A) a (B) b (C) c (D) d (E) e

14. If $y = x^2 \sin 2x$, then $\frac{dy}{dx} =$

- (A) $2x \cos 2x$
(B) $4x \cos 2x$
(C) $2x(\sin 2x + \cos 2x)$
(D) $2x(\sin 2x - x \cos 2x)$
(E) $2x(\sin 2x + x \cos 2x)$
-

Calculus AB

Section I

Part A

15. Let f be the function with derivative given by $f'(x) = x^2 - \frac{2}{x}$. On which of the following intervals is f decreasing?

- (A) $(-\infty, -1]$ only
- (B) $(-\infty, 0)$
- (C) $[-1, 0)$ only
- (D) $(0, \sqrt[3]{2}]$
- (E) $[\sqrt[3]{2}, \infty)$

16. If the line tangent to the graph of the function f at the point $(1, 7)$ passes through the point $(-2, -2)$, then $f'(1)$ is

- (A) -5
- (B) 1
- (C) 3
- (D) 7
- (E) undefined

Unauthorized
Copying
Prohibited
© 2007
College Board
All rights reserved.
www.collegeboard.com

17. Let f be the function given by $f(x) = 2xe^x$. The graph of f is concave down when
- (A) $x < -2$ (B) $x > -2$ (C) $x < -1$ (D) $x > -1$ (E) $x < 0$

x	-4	-3	-2	-1	0	1	2	3	4
$g'(x)$	2	3	0	-3	-2	-1	0	3	2

18. The derivative g' of a function g is continuous and has exactly two zeros. Selected values of g' are given in the table above. If the domain of g is the set of all real numbers, then g is decreasing on which of the following intervals?
- (A) $-2 \leq x \leq 2$ only
(B) $-1 \leq x \leq 1$ only
(C) $x \geq -2$
(D) $x \geq 2$ only
(E) $x \leq -2$ or $x \geq 2$
-


19. A curve has slope $2x + 3$ at each point (x, y) on the curve. Which of the following is an equation for this curve if it passes through the point $(1, 2)$?

- (A) $y = 5x - 3$
- (B) $y = x^2 + 1$
- (C) $y = x^2 + 3x$
- (D) $y = x^2 + 3x - 2$
- (E) $y = 2x^2 + 3x - 3$


$$f(x) = \begin{cases} x + 2 & \text{if } x \leq 3 \\ 4x - 7 & \text{if } x > 3 \end{cases}$$

20. Let f be the function given above. Which of the following statements are true about f ?

- I. $\lim_{x \rightarrow 3} f(x)$ exists.
 - II. f is continuous at $x = 3$.
 - III. f is differentiable at $x = 3$.
- (A) None
 - (B) I only
 - (C) II only
 - (D) I and II only
 - (E) I, II, and III



21. The second derivative of the function f is given by $f''(x) = x(x - a)(x - b)^2$. The graph of f'' is shown above. For what values of x does the graph of f have a point of inflection?
- (A) 0 and a only (B) 0 and m only (C) b and j only (D) 0, a , and b (E) b , j , and k



22. The graph of f' , the derivative of f , is the line shown in the figure above. If $f(0) = 5$, then $f(1) =$
- (A) 0 (B) 3 (C) 6 (D) 8 (E) 11

23. $\frac{d}{dx} \left(\int_0^{x^2} \sin(t^3) dt \right) =$

- (A) $-\cos(x^6)$ (B) $\sin(x^3)$ (C) $\sin(x^6)$ (D) $2x \sin(x^3)$ (E) $2x \sin(x^6)$

24. Let f be the function defined by $f(x) = 4x^3 - 5x + 3$. Which of the following is an equation of the line tangent to the graph of f at the point where $x = -1$?

(A) $y = 7x - 3$

(B) $y = 7x + 7$

(C) $y = 7x + 11$

(D) $y = -5x - 1$

(E) $y = -5x - 5$

25. A particle moves along the x -axis so that at time $t \geq 0$ its position is given by $x(t) = 2t^3 - 21t^2 + 72t - 53$. At what time t is the particle at rest?

(A) $t = 1$ only

(B) $t = 3$ only

(C) $t = \frac{7}{2}$ only

(D) $t = 3$ and $t = \frac{7}{2}$

(E) $t = 3$ and $t = 4$

26. What is the slope of the line tangent to the curve $3y^2 - 2x^2 = 6 - 2xy$ at the point $(3, 2)$?

- (A) 0 (B) $\frac{4}{9}$ (C) $\frac{7}{9}$ (D) $\frac{6}{7}$ (E) $\frac{5}{3}$

27. Let f be the function defined by $f(x) = x^3 + x$. If $g(x) = f^{-1}(x)$ and $g(2) = 1$, what is the value of $g'(2)$?

- (A) $\frac{1}{13}$ (B) $\frac{1}{4}$ (C) $\frac{7}{4}$ (D) 4 (E) 13

Section I

Calculus

Part A


28. Let g be a twice-differentiable function with $g'(x) > 0$ and $g''(x) > 0$ for all real numbers x , such that $g(4) = 12$ and $g(5) = 18$. Of the following, which is a possible value for $g(6)$?
- (A) 15 (B) 18 (C) 21 (D) 24 (E) 27
-

END OF PART A OF SECTION I

CALC all after this point.

76. A particle moves along the x -axis so that at any time $t \geq 0$, its velocity is given by $v(t) = 3 + 4.1 \cos(0.9t)$. What is the acceleration of the particle at time $t = 4$?

(A) -2.016 (B) -0.677 (C) 1.633 (D) 1.814 (E) 2.978



77. The regions A , B , and C in the figure above are bounded by the graph of the function f and the x -axis. If the area of each region is 2, what is the value of $\int_{-3}^3 (f(x) + 1) dx$?




(A) -2 (B) -1 (C) 4 (D) 7 (E) 12

78 *22*

The radius of a circle is increasing at a constant rate of 0.2 meters per second. What is the rate of increase in the area of the circle at the instant when the circumference of the circle is 20π meters?

- (A) $0.04\pi \text{ m}^2/\text{sec}$
- (B) $0.4\pi \text{ m}^2/\text{sec}$
- (C) $4\pi \text{ m}^2/\text{sec}$
- (D) $20\pi \text{ m}^2/\text{sec}$
- (E) $100\pi \text{ m}^2/\text{sec}$

79. For which of the following does $\lim_{x \rightarrow 4} f(x)$ exist?



- (A) I only
- (B) II only
- (C) III only
- (D) I and II only
- (E) I and III only

Used Exam Excerpt

80. The function f is continuous for $-2 \leq x \leq 1$ and differentiable for $-2 < x < 1$. If $f(-2) = -5$ and $f(1) = 4$, which of the following statements could be false?
- (A) There exists c , where $-2 < c < 1$, such that $f(c) = 0$.
 - (B) There exists c , where $-2 < c < 1$, such that $f'(c) = 0$.
 - (C) There exists c , where $-2 < c < 1$, such that $f(c) = 3$.
 - (D) There exists c , where $-2 < c < 1$, such that $f'(c) = 3$.
 - (E) There exists c , where $-2 \leq c \leq 1$, such that $f(c) \geq f(x)$ for all x on the closed interval $-2 \leq x \leq 1$.

-
81. Let f be the function with derivative given by $f'(x) = \sin(x^2 + 1)$. How many relative extrema does f have on the interval $2 < x < 4$?

(A) One (B) Two (C) Three (D) Four (E) Five

82. The rate of change of the altitude of a hot-air balloon is given by $r(t) = t^3 - 4t^2 + 6$ for $0 \leq t \leq 8$. Which of the following expressions gives the change in altitude of the balloon during the time the altitude is decreasing?

(A) $\int_{1.572}^{3.514} r(t) dt$

(B) $\int_0^8 r(t) dt$

(C) $\int_0^{2.667} r(t) dt$

(D) $\int_{1.572}^{3.514} r'(t) dt$

(E) $\int_0^{2.667} r'(t) dt$






83. The velocity, in ft/sec, of a particle moving along the x -axis is given by the function $v(t) = e^t + te^t$. What is the average velocity of the particle from time $t = 0$ to time $t = 3$?

- (A) 20.086 ft/sec
- (B) 26.447 ft/sec
- (C) 32.809 ft/sec
- (D) 40.671 ft/sec
- (E) 79.342 ft/sec

84. A pizza, heated to a temperature of 350 degrees Fahrenheit ($^{\circ}\text{F}$), is taken out of an oven and placed in a 75°F room at time $t = 0$ minutes. The temperature of the pizza is changing at a rate of $-110e^{-0.4t}$ degrees Fahrenheit per minute. To the nearest degree, what is the temperature of the pizza at time $t = 5$ minutes?






- (A) 112°F (B) 119°F (C) 147°F (D) 238°F (E) 335°F
-

85. If a trapezoidal sum overapproximates $\int_0^4 f(x) dx$, and a right Riemann sum underapproximates $\int_0^4 f(x) dx$, which of the following could be the graph of $y = f(x)$?



88. On the closed interval $[2, 4]$, which of the following could be the graph of a function f with the property that

$$\frac{1}{4-2} \int_2^4 f(t) dt = 1?$$



AP Calculus AB Exam

R2

89. Let f be a differentiable function with $f(2) = 3$ and $f'(2) = -5$, and let g be the function defined by $g(x) = xf(x)$. Which of the following is an equation of the line tangent to the graph of g at the point where $x = 2$?

- (A) $y = 3x$
 (B) $y - 3 = -5(x - 2)$
 (C) $y - 6 = -5(x - 2)$
 (D) $y - 6 = -7(x - 2)$
 (E) $y - 6 = -10(x - 2)$

90. For all x in the closed interval $[2, 5]$, the function f has a positive first derivative and a negative second derivative. Which of the following could be a table of values for f ?

(A)

x	$f(x)$
2	7
3	9
4	12
5	16

(B)

x	$f(x)$
2	7
3	11
4	14
5	16

(C)

x	$f(x)$
2	16
3	12
4	9
5	7

(D)

x	$f(x)$
2	16
3	14
4	11
5	7

(E)

x	$f(x)$
2	16
3	13
4	10
5	7

91. A particle moves along the x -axis so that at any time $t > 0$, its acceleration is given by $a(t) = \ln(1 + 2^t)$. If the velocity of the particle is 2 at time $t = 1$, then the velocity of the particle at time $t = 2$ is

- (A) 0.462 (B) 1.609 (C) 2.555 (D) 2.886 (E) 3.346

92. Let g be the function given by $g(x) = \int_0^x \sin(t^2) dt$ for $-1 \leq x \leq 3$. On which of the following intervals is g decreasing?

- (A) $-1 \leq x \leq 0$
(B) $0 \leq x \leq 1.772$
(C) $1.253 \leq x \leq 2.171$
(D) $1.772 \leq x \leq 2.507$
(E) $2.802 \leq x \leq 3$

END OF SECTION I

AFTER TIME HAS BEEN CALLED, TURN TO THE NEXT PAGE AND
ANSWER QUESTIONS 93-96.